

Lead Toxicity Overview

- **There is NO safe level of lead in the blood of a child.**
- Exposure to lead can seriously harm a child's health and cause well-documented adverse effects such as:
 - Damage to the brain and nervous system
 - Slowed growth and development
 - Learning and behavior problems and underperformance in school
 - Lower IQ and decreased ability to pay attention
 - Hearing and speech problems
- Lead is particularly dangerous because once in a person's system, it is distributed throughout the body, causing harm wherever it lands. In the bloodstream, it can damage red blood cells and limit their ability to carry oxygen to organs and tissues, causing anemia. Most lead ends up in the bone where it interferes with the production of red blood cells and the absorption of calcium. The effects of long-term lead toxicity include damage to the nervous system, kidneys, and hearing, poor muscle coordination, and decreased bone and muscle growth.
- **The American Academy of Pediatrics recommends:**
 - Physicians continue to provide anticipatory guidance to parents in an effort to prevent lead exposure
 - Pediatricians should increase their efforts to assess children at risk for lead exposure to find those with elevated BLLs
- **Populations at higher risk of lead exposure include:**
 - Children from low-income households,
 - Immigrant and refugee children from less developed countries
 - Some African American children, and those from other races and ethnicities, due to poor housing stock
 - Those living in housing built before 1978
 - Children younger than age 6 (hand to mouth activity)

However, children from all races, ethnicities and socioeconomic backgrounds can be at risk of exposure.

- All homes built prior to 1978 have a risk for lead exposure, but homes built before 1950 carry a greater risk due to a higher content of lead used in paint before that time. Age and condition of housing, **not geographic location**, are the **best predictors** for the presence of hazards related to lead-based paint. Besides lead-based paint, secondary sources of lead continue to emerge (e.g., children's jewelry, imported herbal supplements, and food products) and are not limited to high-risk populations.

Indiana Lead Testing Requirements

- **Children Enrolled in Medicaid:** Indiana statute requires that ALL children enrolled in Medicaid are to receive a blood lead level test at **12 and 24 months** of age; ANY child enrolled in Medicaid between 36 and 72 months with no record of a previous blood lead test must also be tested.
- **Additional testing should be strongly considered if children:**
 - Live in a house built before 1978 or a home that was recently repaired or renovated.
 - Are enrolled in Women, Infant and Children Supplemental Nutrition Program (WIC) or Head Start.
 - Have any of the risk factors in the Verbal Risk Assessment (below).
 - Are adopted outside the United States, in foster care, or are immigrants.
 - Have a known history of lead exposure after the age of 2 years old.
 - Have a sibling or a playmate with lead toxicity.
 - Have parents who request testing.
 - Live near a lead-emitting facility.
 - Are exhibiting neurodevelopmental disabilities or conditions such as autism, attention-deficit/hyperactivity disorder, and learning delays.
 - Have a history of ingested non-food items (e.g. jewelry, batteries, toys, etc.) or exhibit pica behavior.

Indiana Lead Testing Recommendations

- **Testing Refugee Populations:** CDC recommends refugee children aged 6 months to 16 years should be tested upon entry into the United States. Testing should be repeated 3 to 6 months after placement in a permanent residence, regardless of initial test results. All pregnant and lactating women and girls should be tested. All newly arrived pregnant or breastfeeding women should be prescribed a prenatal or multivitamin with adequate iron and calcium. Referral to a healthcare provider with expertise in high-risk lead exposure treatment and management may be indicated for EBLLs
- It is recommended that clinicians **assess all children** for the risk of lead exposure from ages 6 months to 6 years **at every well child visit.**

Verbal Risk Assessment:

The **Verbal Risk Assessment** helps health care providers assess a child’s potential level of risk of exposure to lead hazards. It also provides health care providers an opportunity to educate families about lead hazards. **If the answer to any question on the Verbal Risk Assessment is “Yes” or “I don’t know,”** then:

- Test the child for lead toxicity at that time
- Test the child at least annually or if any other risk factor exists. Continue to test until the child turns 6 years old or responses to the assessment change.
- Provide lead toxicity prevention education

* Completion of the assessment does not meet the Medicaid testing requirement.

In the past year, has your child lived in, near, or regularly visited:	Yes	No	I don’t know
A house built before 1978 that has peeling, chipping, or flaking paint?			
A house built before 1978 that has been remodeled within the past 6 months?			
A sibling, cousin, or friend who has been diagnosed or treated for lead toxicity?			
A factory or industrial plant or mine?			
Mexico, India, Middle East, Central America, South America, Africa, or Asia			
In the past year, has your child been around adults who:	Yes	No	I don’t know
Have a job that causes them to have frequent contact with lead (e.g. plumbers, construction, auto repair, metal/battery recycling, welders)			
Have a hobby that causes them to have frequent contact with lead (e.g. hunt, fish, reload bullets, refinish furniture, work with stained glass, jewelry making)			
In the past year, has your child consumed:	Yes	No	I don’t know
Food or beverages from ceramic cookware/dishware or imported pottery?			
Food with spices imported or brought in from another country (such as turmeric)?			
Ayurvedic medicines or home remedies (such as Azarcon, Greta)?			
Dirt or non-food items regularly (more than the typical baby mouthing behavior)?			

Childhood Blood Lead Medical Management Guidelines for Providers in Indiana

General medical evaluation and testing recommendations:

- Perform routine history and assessment of physical and mental development.
- Assess nutrition and risk for iron deficiency.
- Evaluate for lead exposure risks.
- Initial test may be a capillary or venous test. Children with identified risk factors should be retested with a venous sample.

General clinical management recommendations:

- Report the blood lead result to IDOH.
- Discuss result with family and counsel on any identified risk factors.
- Provide prevention education (i.e., lead sources, reducing/eliminating strategies, health education, etc.)
- Counsel on healthy eating, especially iron, calcium, and Vitamin C.
- Communicate patient management information with local health department.
- Consider referral to Supplemental Nutrition Program for Women, Infants, and Children.

Medical Evaluation and Testing Recommendations	Clinical Management
BLL <5 µg/dL	
<p>General medical evaluation recommendations (given above)</p> <p>Who to test?</p> <ul style="list-style-type: none"> • Medicaid recipients at 12 and 24 months, or anytime before 6 years old if not previously screened. • Children in homes built before 1978 or with other risk factors • Children at the request of their parents. 	<p>General clinical management recommendations (given above)</p> <ul style="list-style-type: none"> • When reviewing lab results with family, for reference, the geometric mean blood lead level for children 1-5 years old is less than 2 µg/dL. <p>Chelation is NOT recommended in this blood lead level range.</p>
BLL 5-9.9 µg/dL	
<p>General medical evaluation recommendations, PLUS:</p> <ul style="list-style-type: none"> • Note the child's environmental history. Identify potential sources of exposure and provide preliminary advice on reducing/eliminating them. • Ensure iron sufficiency with laboratory testing and treatment per Pediatric Environmental Health Specialty Unit (PEHSU) and American Academy of Pediatrics guidelines. • Perform structured developmental screening evaluations at periodic health visits as lead effects may manifest over years. • Evaluate risk to household contacts such as siblings and pregnant/lactating women in the home. <p>Monitor BLLs:</p> <ul style="list-style-type: none"> • Retest of BLL done no later than 6 months, recommended within shorter time period to ensure blood lead level decreasing. • If retest result is in another range, follow up as for that range. • If BLLs are stable or decreasing monitor initially with venous BLLs every 3 months and thereafter based on venous BLL trend. If retest result is in another range, follow-up or retest as for that range. 	<p>General clinical management recommendations, PLUS:</p> <ul style="list-style-type: none"> • Assess the child's environmental risk factors, eating habits, housing, and family's social service needs. • Test for iron sufficiency. Consider starting a multivitamin tablet with iron. • Test siblings, other children <6 years old, and household contacts, especially pregnant and lactating women • Include primary/secondary residence and child care facility as part of the investigation. <p>Chelation is NOT recommended in this BLL range.</p>
BLL 10-14.9 µg/dL	
<p>General medical evaluation recommendations, PLUS:</p> <ul style="list-style-type: none"> • Note the child's environmental history. Identify potential sources of exposure and provide preliminary advice on reducing/eliminating them. • Ensure iron and calcium sufficiency with laboratory testing and treatment per Pediatric Environmental Health Specialty Unit (PEHSU) and American Academy of Pediatrics guidelines. • Perform structured developmental screening evaluations at 	<p>General clinical management recommendations, PLUS:</p> <ul style="list-style-type: none"> • Assess the child's environmental risk factors, eating habits, housing, and family's social service needs. • If a past exposure is noted, perform developmental screenings at periodic health visits. Health effects of lead manifest over time. • Test for iron sufficiency. Consider starting a multivitamin tablet with iron. • Test siblings, other children <6 years old, and household

Childhood Blood Lead Medical Management Guidelines for Providers in Indiana

<p>periodic health visits as lead effects may manifest over years.</p> <ul style="list-style-type: none"> Evaluate risk to household contacts such as siblings and pregnant/lactating women in the home. <p>Monitor BLLs:</p> <ul style="list-style-type: none"> Confirmatory BLL draw within 3 months of original. Venous test preferred, capillary acceptable Retest thereafter within 3 months until BLL declines If retest result is in another range, follow up as for that range. If BLLs are stable or decreasing, monitor initially with venous BLLs based on venous BLL trend. If retest result is in another range, follow-up or retest as for that range. 	<p>contacts, especially pregnant and lactating women</p> <ul style="list-style-type: none"> Make referrals to the local social services and assistance (WIC, Head Start, First Steps, Help Me Grow, etc.) office if necessary. Include primary/secondary residence and child care facility as part of the investigation. Environmental investigation and recommendations for remediation services and home visit (education/referrals) by LHD staff will be completed at this BLL. If BLL is persistent or rising, contact the local health department (LHD). <p>Chelation is NOT recommended in this BLL range.</p>
BLL 15-19.9 µg/dL	
<p>General medical evaluation recommendations:</p> <p>Monitor BLLs:</p> <ul style="list-style-type: none"> Confirmatory BLL draw within 3 months of original. Venous test preferred, capillary acceptable Retest thereafter within 3 months until BLL declines If retest result is in another range, follow up as for that range. If BLLs are stable or decreasing, monitor initially with venous BLLs based on venous BLL trend. If retest result is in another range, follow-up or retest as for that range. 	<p>General clinical management recommendations:</p> <p>Chelation is NOT recommended in this BLL range.</p>
BLL 20-44.9 µg/dL	
<p>General medical evaluation recommendations, PLUS:</p> <ul style="list-style-type: none"> Complete diagnostic evaluation, history and physical exam with neurodevelopmental assessment. <p>Monitor BLLs:</p> <ul style="list-style-type: none"> Confirmatory BLL draw within 1 month. Venous test preferred, capillary acceptable Retest within 1 to 3 months, depending on range, to ensure BLL is not rising. Monitor monthly and afterward based on the BLL trend. If retest result is in another range, follow up as for that range. 	<p>General clinical management recommendations, PLUS:</p> <ul style="list-style-type: none"> Additional evaluation such as abdominal x-ray should be considered based on environmental investigation (e.g. pica for paint chips, mouthing behaviors, etc.) Gut decontamination may be considered if leaded foreign bodies visualized on x-ray. <p>Chelation is NOT recommended in this BLL range.</p>
BLL 45-69.9 µg/dL (Urgent Medical Situation)	
<p>General medical evaluation recommendations, PLUS:</p> <ul style="list-style-type: none"> Any treatment for BLLs in this range and greater should be done in consultation with an expert. Contact IN Poison Control (1-800-222-1222) or Region 5 PEHSU (1-866-967-7337) for guidance regarding chelation, diagnostic tests, questions, etc. <p>Monitor BLLs:</p> <ul style="list-style-type: none"> Confirmatory BLL draw within 24 hours. Consider reconfirming BLL, even for venous results. Retesting dependent on BLL range. Retest, venous only, 1 month after chelation therapy is completed. If confirmed in this range, monitor BLLs during chelation. Retest every 2 to 4 weeks (or more based on most recent BLLs). Modify treatment guidelines if BLL remains elevated. <p>Monitor frequently until BLL declines.</p>	<p>General clinical management recommendations, PLUS:</p> <ul style="list-style-type: none"> Evaluate whether hospitalization is needed to reduce lead exposure (Managed with assistance of experienced provider). Notify LHD if admitted or chelation administered. <p><u>Consider chelation therapy:</u></p> <ul style="list-style-type: none"> Consult with a provider experienced in managing chelation therapy. Chelation may be conducted at child's home if no lead hazards are present. If hazards are present, child must be admitted to hospital to receive chelation therapy. Consider bowel decontamination as an adjunct to chelation if abdominal X-ray indicates enteral lead is present. Succimer can be prescribed. A minimum of two weeks between courses is recommended,

Childhood Blood Lead Medical Management Guidelines for Providers in Indiana

	<p>unless more prompt treatment is indicated.</p> <ul style="list-style-type: none"> • Discontinue iron supplements. • Monitor for anemia and neutropenia. <p><u>Post-Chelation Therapy Guidelines:</u></p> <ul style="list-style-type: none"> • Repeat venous lead test in 1 to 3 weeks after hospital discharge. • Repeat venous lead test every two weeks for 6 to 8 weeks after hospital discharge. • Monitor lead level closely for 4 to 6 months after chelation. If the lead level “rebounds” to pre-treatment levels, consider repeat chelation therapy. • Minimum of two-week intervals is needed between chelation courses.
BLL ≥ 70 µg /dL (Medical Emergency)	
<p>General medical evaluation recommendations:</p> <p>Monitor BLLs:</p> <ul style="list-style-type: none"> • Confirmatory BLL draw done immediately, emergency lab test, considered Medical Emergency. Venous test is preferred. • Retesting dependent on BLL range. Retest, venous only, 1 month after chelation is completed • Retest within 1 week to 1 month to ensure BLL is not rising. • Monitor monthly and afterward based on the BLL trend. If retest result is in another range, follow up as for that range. • Any treatment for BLLs in this range should be done in consultation with an expert. • Refer to CDC and American Academy of Pediatrics recommendations related to chelation management. 	<p>General clinical management recommendations, PLUS:</p> <p>Follow chelation therapy and post-chelation therapy guidelines listed above.</p>

Resources:

1. CDC: <https://www.cdc.gov/nceh/lead/>
2. Environmental Health Specialty Units (PEHSUs): <https://childrensenviron.uic.edu/>
3. AAPs Bright Futures Guidelines: <https://brightfutures.aap.org/Pages/default.aspx>
4. Indiana Department of Health, Lead and Healthy Homes Division: <https://www.in.gov/health/lead-and-healthy-homes-division/>
5. Indiana Poison Control: 1-800-222-1222