ABATACEPT IV

Products Affected

• ORENCIA (WITH MALTOSE)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	RA, PJIA, PSA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. AGVHD: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ABATACEPT SQ

Products Affected

- ORENCIA
- ORENCIA CLICKJECT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

ABEMACICLIB

Products Affected

VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ABIRATERONE

Products Affected

• abiraterone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER, METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ABIRATERONE SUBMICRONIZED

Products Affected

· YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ACALABRUTINIB

Products Affected

- CALQUENCECALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ADAGRASIB

Products Affected

KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ADALIMUMAB

Products Affected

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UVEITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)

- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OPTHALMOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIEN E, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, PJIA, PSA, AS, PSO, HIDRADENITIS SUPPURATIVA, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AFATINIB

Products Affected

• GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ALECTINIB

Products Affected

· ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ALPELISIB

Products Affected

 PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AMBRISENTAN

Products Affected

• ambrisentan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL: PATIENT DOES NOT HAVE IDIOPATHIC PULMONARY FIBROSIS. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR 2) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AMIVANTAMAB-VMJW

Products Affected

RYBREVANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APALUTAMIDE

Products Affected

• ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): (1) HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): (1) RECEIVED A BILATERAL ORCHIECTOMY, (2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR (3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE ANALOG. RENEWAL: DIAGNOSIS OF NMCRPC OR MCSPC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APOMORPHINE

Products Affected

• apomorphine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PD. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WHILE ON THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APOMORPHINE - SL

Products Affected

 KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF KYNMOBI.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

APREMILAST

Products Affected

- OTEZLA
- OTEZLA STARTER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: MILD PLAQUE PSORIASIS (PSO): ONE OF THE FOLLOWING: 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: 1) PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR 2) PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC AGENT (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) AND ONE CONVENTIONAL TOPICAL AGENT (E.G., PUVA, UVB, TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. BEHCETS DISEASE: 1) PATIENT HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: PSA, PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ASCIMINIB

Products Affected

SCEMBLIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ASFOTASE

Products Affected

• STRENSIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6
	MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2)
	POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE
	PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS
	CONFIRMED BY GENETIC TESTING OR TWO OF THE
	FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP)
	LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT
	AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE
	(PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN
	THE PREVIOUS WEEK, (C) URINE
	PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF
	NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC
	EVIDENCE OF HPP, (E) AT LEAST TWO OF THE
	FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II)
	CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH
	RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV)
	HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V)
	NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM
	CALCIUM, (VI) HISTORY OR PRESENCE OF NON-
	TRAUMATIC POSTNATAL FRACTURE AND DELAYED
	FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS
	OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE
	FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED
	BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A)
	SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE
	FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS
	AND NO VITAMIN B6 SUPPLEMENTATION IN THE
	PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF
	NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC
	EVIDENCE OF HPP, (E) AT LEAST TWO OF THE
	FOLLOWING: (I) RACHITIC DEFORMITIES, (II)
	PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5
	YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH
	RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV)

PA Criteria	Criteria Details
	HISTORY OR PRESENCE OF NON-TRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3) NOT HAVE A TREATABLE FORM OF RICKETS. RENEWAL: ALL INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ATOGEPANT

Products Affected

• QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AVAPRITINIB

Products Affected

AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AVATROMBOPAG

Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC LIVER DISEASE (CLD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, GASTROENTEROLOGIST, HEPATOLOGIST, IMMUNOLOGIST, ENDOCRINOLOGIST, OR A SURGEON. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	CLD: 1 MONTH. CHRONIC ITP: INITIAL: 2 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: CLD: 1) PLANNED PROCEDURE 10 TO 13 DAYS AFTER INITIATION OF DOPTELET, AND 2) NOT RECEIVING OTHER THROMBOPOIETIN RECEPTOR AGONISTS (E.G., ROMIPLOSTIM, ELTROMBOPAG, ETC.). CHRONIC ITP: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS OR IMMUNOGLOBULINS OR INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CHRONIC ITP: PATIENT HAD A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AXITINIB

Products Affected

• INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AZACITIDINE

Products Affected

ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

AZTREONAM

Products Affected

CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	7 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BARICITINIB

Products Affected

OLUMIANT

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SEVERE ALOPECIA AREATA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	RA, SEVERE ALOPECIA AREATA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO ANY TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ. SEVERE ALOPECIA AREATA: 1) AT LEAST 50 PERCENT SCALP HAIR LOSS AS MEASURED BY THE SEVERITY OF ALOPECIA TOOL (SALT) FOR MORE THAN 6 MONTHS, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR SEVERE ALOPECIA AREATA OR OTHER JAK INHIBITORS FOR ANY INDICATION. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR SEVERE ALOPECIA AREATA OR OTHER JAK INHIBITORS FOR ANY INDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

BEDAQUILINE

Products Affected

SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 WEEKS
Other Criteria	PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS (MDR-TB): SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MDR-TB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BELIMUMAB

Products Affected

• BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BELUMOSUDIL

Products Affected

REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BELZUTIFAN

Products Affected

• WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BENDAMUSTINE

- bendamustine intravenous recon soln
- BENDAMUSTINE INTRAVENOUS SOLUTION
- BENDEKA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BENRALIZUMAB

- FASENRA
- FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: 4 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEVI FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BETAINE

Products Affected

• betaine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEVACIZUMAB-ADCD

Products Affected

VEGZELMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEVACIZUMAB-AWWB

Products Affected

MVASI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEVACIZUMAB-BVZR

Products Affected

ZIRABEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BEXAROTENE

Products Affected

• bexarotene

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BINIMETINIB

Products Affected

MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BORTEZOMIB

- bortezomib injection recon soln 1 mg, 2.5 VELCADE mg
- BORTEZOMIB INTRAVENOUS RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BOSENTAN

- TRACLEER ORAL TABLET
- TRACLEER ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE. RENEWAL: 1) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE, 2) AGES 3 TO 17 YEARS OF AGE: A) DEMONSTRATED IMPROVEMENT IN PVR, OR B) REMAINED STABLE OR SHOWN IMPROVEMENT IN EXERCISE ABILITY, 3) AGES 18 YEARS OR OLDER: A) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR B) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BOSUTINIB

Products Affected

• BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

BRIGATINIB

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS, DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

C1 ESTERASE INHIBITOR-CINRYZE

Products Affected

CINRYZE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

C1 ESTERASE INHIBITOR-HAEGARDA

Products Affected

• HAEGARDA SUBCUTANEOUS RECON SOLN 2,000 UNIT, 3,000 UNIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CABOZANTINIB

Products Affected

 COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CABOZANTINIB S-MALATE - CABOMETYX

Products Affected

• CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CANNABIDIOL

Products Affected

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DS, LGS, TSC: CONFIRMATION OF DIAGNOSIS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CAPLACIZUMAB YHDP

Products Affected

CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACQUIRED THROMBOTIC THROMBOCYTOPENIA PURPURA (ATTP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	ATTP: CABLIVI WAS PREVIOUSLY INITIATED AS PART OF THE FDA APPROVED TREATMENT REGIMEN IN COMBINATION WITH PLASMA EXCHANGE AND IMMUNOSUPPRESSIVE THERAPY WITHIN AN INPATIENT SETTING. THE PATIENT HAS NOT EXPERIENCED MORE THAN TWO RECURRENCES OF ATTP WHILE ON CABLIVI THERAPY (I.E., NEW DROP IN PLATELET COUNT REQUIRING REPEAT PLASMA EXCHANGE DURING 30 DAYS POST-PLASMA EXCHANGE THERAPY [PEX] AND UP TO 28 DAYS OF EXTENDED THERAPY).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CAPMATINIB

Products Affected

TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CARGLUMIC ACID

Products Affected

• carglumic acid

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ACUTE, CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY THE PRESENCE OF ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMING MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY THE PRESENCE OF ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMING MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.
Other Criteria	RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CERITINIB

Products Affected

ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CERTOLIZUMAB PEGOL

- CIMZIA
- CIMZIA POWDER FOR RECONST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI, RINVOQ. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, RINVOQ. PATIENTS WHO ARE PREGNANT, BREASTFEEDING, OR TRYING TO BECOME PREGNANT ARE EXCLUDED FROM STEP CRITERIA FOR ALL INDICATIONS. RENEWAL: RA, PSA, AS, PSO, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CETUXIMAB

Products Affected

• ERBITUX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CLADRIBINE

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 48 WEEKS.
Other Criteria	MS: INITIAL: HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF 2 CYCLES IN EACH). RENEWAL: 1) HAS DEMONSTRATED CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE, 2) DOES NOT HAVE LYMPHOPENIA, AND 3) HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF 2 CYCLES IN EACH).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CLOBAZAM-SYMPAZAN

Products Affected

SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	LGS: 1) PATIENT IS UNABLE TO TAKE TABLETS OR SUSPENSION, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY CLOBAZAM AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

COBIMETINIB

Products Affected

· COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

COLCHICINE

Products Affected

• colchicine oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PROPHYLAXIS OF GOUT FLARES: 16 YEARS AND OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PROPHYLAXIS OF GOUT FLARES: TRIAL OF OR CONTRAINDICATION TO COLCHICINE CAPSULES (MITIGARE) IF AGE 18 YEARS OR OLDER.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CORTICOTROPIN

- ACTHAR
- · CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.
Coverage Duration	INFANTILE SPASMS AND MS: 28 DAYS. OTHER FDA APPROVED INDICATIONS: INITIAL AND RENEWAL: 28 DAYS
Other Criteria	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: TRIAL OF OR CONTRAINDICATION TO A STANDARD OF CARE THERAPY. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: 1) DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS, AND 2) CONTINUES TO POSSESS CONTRAINDICATION TO IV CORTICOSTEROIDS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MAPD PLAN.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

CRIZOTINIB

Products Affected

XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

CYSTEAMINE HYDROCHLORIDE

Products Affected

· CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DABRAFENIB

Products Affected

• TAFINLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DABRAFENIB SUSPENSION

Products Affected

 TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNABLE TO SWALLOW TAFINILAR CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DACOMITINIB

Products Affected

VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DALFAMPRIDINE

Products Affected

• dalfampridine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DAROLUTAMIDE

Products Affected

• NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): PATIENT HAS HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS) AND ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. METASTATIC HORMONE-SENSITIVE PROSTATE CANCER (MHSPC): 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC: NO ADDITONAL CRITERIA REQUIRED. MHSPC: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DASATINIB

Products Affected

 SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED Ph+ CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SPRYCEL IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DECITABINE/CEDAZURIDINE

Products Affected

· INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DEFERASIROX

Products Affected

• deferasirox

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G DRY WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G DRY WEIGHT OR GREATER.
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL FOR ALL INDICATIONS: FORMULARY VERSION OF DEFERASIROX SPRINKLE: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX TABLET OR TABLET FOR ORAL SUSPENSION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

DEFERIPRONE

Products Affected

- deferiprone FERRIPROX (2 TIMES A DAY) FERRIPROX ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TRANSFUSIONAL IRON OVERLOAD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: TRANSFUSIONAL IRON OVERLOAD DUE TO THALASSEMIA SYNDROMES: 1) TRIAL OF, CONTRAINDICATION, INTOLERABLE TOXICITIES, OR CLINICALLY SIGNIFICANT ADVERSE EFFECTS TO ONE OF THE FOLLOWING: FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE, OR 2) CURRENT CHELATION THERAPY (I.E., FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE) IS INADEQUATE. TRANSFUSIONAL IRON OVERLOAD DUE TO SICKLE CELL DISEASE OR OTHER ANEMIAS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE. RENEWAL (ALL INDICATIONS): SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 500MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

DENOSUMAB-XGEVA

Products Affected

XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DEUTETRABENAZINE

Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG,
- 24 MG, 6 MG
- AUSTEDO XR TITRATION KT(WK1-4)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TARDIVE DYSKINESIA: PATIENT HAS A HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DEXMEDETOMIDINE

Products Affected

• IGALMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACUTE AGITATION: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.
Coverage Duration	INITIAL: 1 MO. RENEWAL: SEE OTHER CRITERIA.
Other Criteria	ACUTE AGITATION: INITIAL: TRIAL AND FAILURE OF OR CONTRAINDICATION TO TWO ANTIPSYCHOTICS. RENEWAL: 1) MAINTENANCE THERAPY FOR UNDERLYING PSYCHIATRIC DISORDER IS CURRENTLY ADJUSTED/OPTIMIZED TO REDUCE/ELIMINATE THE NEED FOR CONTINUED PRN MEDICATIONS: 3 MONTHS, OR 2) ATTEMPTS TO ADJUST MEDICATIONS HAVE BEEN EXHAUSTED AND PHYSICIAN DETERMINED THAT CHRONIC PRN MEDICATIONS ARE REQUIRED FOR THE CONTINUED SAFETY OF PATIENT OR CAREGIVERS: 6 MONTHS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DICLOFENAC GEL

Products Affected

• diclofenac sodium topical gel 3 %

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DICLOFENAC TOPICAL SOLUTION

Products Affected

• diclofenac sodium topical solution in metered-dose pump

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DIMETHYL FUMARATE

Products Affected

• dimethyl fumarate oral capsule,delayed release(drlec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DIROXIMEL FUMARATE

Products Affected

VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DOSTARLIMAB-GXLY

Products Affected

• JEMPERLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DRONABINOL

Products Affected

• dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DROXIDOPA

Products Affected

• droxidopa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION.
Age Restrictions	
Prescriber Restrictions	NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
Other Criteria	NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

DUPILUMAB

Products Affected

- DUPIXENT PEN
- DUPIXENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY.
Age Restrictions	
Prescriber Restrictions	INITIAL: AD, PN: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST.
Coverage Duration	INITIAL: AD, CRSWNP, EOE, PN: 6 MOS, ASTHMA: 4 MOS. RENEWAL: ALL INDICATIONS: 12 MOS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: AD: 1) AD COVERING AT LEAST 10 PERCENT OF
	BODY SURFACE AREA OR AD AFFECTING THE FACE,
	HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS
	AREAS, 2) INTRACTABLE PRURITUS OR
	CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3)
	TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL
	(CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4
	INHIBITOR, OR JAK INHIBITOR), AND 4) NO
	CONCURRENT USE WITH OTHER SYSTEMIC
	BIOLOGIC/JAK INHIBITOR FOR AD. ASTHMA: 1)
	CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE
	OR MAXIMALLY-TOLERATED DOSE OF AN INHALED
	CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE
	MEDICATION, 2) ONE ASTHMA EXACERBATION
	REQUIRING SYSTEMIC CORTICOSTEROID BURST
	LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS,
	OR ONE SERIOUS EXACERBATION REQUIRING
	HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12
	MONTHS, OR POOR SYMPTOM CONTROL DESPITE
	CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE
	OF THE FOLLOWING WITHIN THE PAST 4 WEEKS:
	DAYTIME ASTHMA SYMPTOMS MORE THAN
	TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA,
	SABA RELIEVER FOR SYMPTOMS MORE THAN
	TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO
	ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR OR
	OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA.
	CRSWNP: 1) EVIDENCE OF NASAL POLYPS BY DIRECT
	EXAMINATION, ENDOSCOPY OR SINUS CT SCAN, 2)
	INADEQUATELY CONTROLLED DISEASE AS
	DETERMINED BY USE OF SYSTEMIC STEROIDS IN THE
	PAST 2 YEARS OR ENDOSCOPIC SINUS SURGERY, AND 3) A
	90 DAY TRIAL OF ONE TOPICAL NASAL
	CORTICOSTEROID. PN: 1) CHRONIC PRURITIS (ITCH MORE

PA Criteria	Criteria Details
	THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, AND
	HISTORY OR SIGN OF A PROLONGED SCRATCHING
	BEHAVIOR, 2) TRIAL OF OR CONTRAINDICATION TO ONE
	TOPICAL (CORTICOSTEROID OR CALCIPOTRIOL).
	RENEWAL: AD: 1) IMPROVEMENT WHILE ON THERAPY,
	AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC
	BIOLOGIC/JAK INHIBITOR FOR AD. CRSWNP, EOE:
	IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) NO
	CONCURRENT USE WITH XOLAIR, OR OTHER ANTI-IL5
	BIOLOGICS FOR ASTHMA, 2) CONTINUED USE OF ICS AND
	ONE OTHER MAINTENANCE MEDICATION, AND 3)
	CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION
	IN ASTHMA EXACERBATIONS FROM BASELINE, (B)
	DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C)
	INCREASE IN PERCENT PREDICTED FEV1 FROM
	PRETREATMENT BASELINE, OR (D) REDUCTION IN
	SEVERITY OR FREQUENCY OF ASTHMA-RELATED
	SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF
	PRURITIS OR PRURIGINOUS LESIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B	No
Prerequisite	

DUVELISIB

Products Affected

COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELACESTRANT

Products Affected

• ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELAGOLIX SODIUM

Products Affected

 ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO A NSAID AND PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

ELEXACAFTOR-TEZACAFTOR-IVACAFTOR

Products Affected

• TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELIGLUSTAT

Products Affected

· CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELRANATAMAB-BCMM

Products Affected

- ELREXFIO 44 MG/1.1 ML VIAL
- ELREXFIO SUBCUTANEOUS SOLUTION 40 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RELAPSED OR REFRACTORY MULTIPLE MYELOMA: RENEWAL: 1) HAS RECEIVED AT LEAST 24 WEEKS OF TREATMENT WITH ELREXFIO, AND 2) HAS RESPONDED TO TREATMENT (PARTIAL RESPONSE OR BETTER), AND HAS MAINTAINED THIS RESPONSE FOR AT LEAST 2 MONTHS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ELTROMBOPAG

Products Affected

- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	ITP: INITIAL: 2 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
Other Criteria	INITIAL: PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA PURPURA (ITP): TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. ALL INDICATIONS: APPROVAL FOR PROMACTA ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF PROMACTA TABLETS OR PATIENT IS UNABLE TO TAKE TABLET FORMULATION. RENEWAL: ITP: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EMAPALUMAB-LZSG

Products Affected

GAMIFANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH): INITIAL: 1) A GENETIC TEST IDENTIFYING HLH-ASSOCIATED GENE MUTATION (E.G., PRF1, UNC13D), OR 2) HAS AT LEAST FIVE OF THE FOLLOWING EIGHT DIAGNOSTIC CRITERIA FOR HLH: (A) FEVER, (B) SPLENOMEGALY, (C) CYTOPENIAS (AFFECTING AT LEAST 2 OF 3 CELL LINEAGES), (D) HYPERTRIGLYCERIDEMIA OR HYPOFIBRINOGENEMIA, (E) HEMOPHAGOCYTOSIS IN BONE MARROW OR SPLEEN OR LYMPH NODES AND NO EVIDENCE OF MALIGNANCY, (F) LOW OR ABSENT NATURAL KILLER-CELL ACTIVITY, (G) FERRITIN LEVEL OF 500 MCG/L OR GREATER, (H) SOLUBLE CD25 LEVEL OF 2,400 U/ML OR GREATER.
Age Restrictions	
Prescriber Restrictions	HLH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN IMMUNOLOGIST, HEMATOLOGIST, OR ONCOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 8 WEEKS.

PA Criteria	Criteria Details
Other Criteria	HLH: INITIAL: 1) CONCURRENT THERAPY WITH DEXAMETHASONE, AND 2) ONE OF THE FOLLOWING: (A) HAS REFRACTORY, RECURRENT, OR PROGRESSIVE DISEASE, OR (B) HAD A TRIAL OF OR INTOLERANCE TO CONVENTIONAL HLH THERAPY (I.E., CHEMOTHERAPY, STEROIDS, IMMUNOTHERAPY). RENEWAL: 1) HAS NOT RECEIVED SUCCESSFUL HEMATOPOIETIC STEM CELL TRANSPLANTATION, AND 2) DEMONSTRATED IMPROVED IMMUNE SYSTEM RESPONSE FROM BASELINE (E.G., RESOLUTION OF FEVER, DECREASED SPLENOMEGALY, IMPROVEMENT IN CNS SYMPTOMS, IMPROVED CBC, INCREASED FIBRINOGEN LEVELS, REDUCED D-DIMER, REDUCED FERRITIN, REDUCED SOLUBLE CD25 LEVELS.)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENASIDENIB

Products Affected

• IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENCORAFENIB

Products Affected

• BRAFTOVI ORAL CAPSULE 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENTRECTINIB

Products Affected

 ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ENZALUTAMIDE

Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: CASTRATION-RESISTANT PROSTATE CANCER (CRPC) THAT IS NOT METASTATIC: PATIENT HAS HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). CRPC (INCLUDES NON-METASTATIC AND METASTATIC) OR METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: DIAGNOSIS OF CRPC (INCLUDES NON-METASTATIC AND METASTATIC) OR MCSPC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EPCORITAMAB-BYSP

Products Affected

EPKINLY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EPOETIN ALFA-EPBX

Products Affected

 RETACRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/2 ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL OF LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL LESS THAN 13G/DL. RENEWAL: CKD: 1) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR 2) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS. ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. CANCER CHEMOTHERAPY: 1) HEMOGLOBIN LEVEL OF LESS THAN 10 G/DL, OR 2) THE HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.
Other Criteria	RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ERDAFITINIB

Products Affected

 BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ERLOTINIB

Products Affected

• erlotinib oral tablet 100 mg, 150 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ESKETAMINE

Products Affected

• SPRAVATO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.
Coverage Duration	INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.
Other Criteria	INITIAL: TRD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, 2) NO ACTIVE SUBSTANCE ABUSE, AND 3) ADEQUATE TRIAL (AT LEAST 4 WEEKS) OF AT LEAST TWO ANTIDEPRESSANT AGENTS FROM DIFFERENT CLASSES THAT ARE INDICATED FOR DEPRESSION. MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ETANERCEPT

Products Affected

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), PSORIATIC ARTHRITIS (PSA): 18 YEARS OR OLDER.
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), AS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE: IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: RA, PJIA, PSA, AS, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

EVEROLIMUS

Products Affected

- everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg
- everolimus (antineoplastic) oral tablet for suspension

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FECAL MICROBIOTA CAPSULE

Products Affected

VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	30 DAYS
Other Criteria	CLOSTRIDIOIDES DIFFICILE INFECTION (CDI): 1) HAS NOT PREVIOUSLY RECEIVED VOWST: COMPLETION OF ANTIBIOTIC TREATMENT FOR RECURRENT CDI (AT LEAST 3 CDI EPISODES), OR 2) PREVIOUSLY RECEIVED VOWST: (A) TREATMENT FAILURE (DEFINED AS THE PRESENCE OF CDI DIARRHEA WITHIN 8 WEEKS OF FIRST DOSE OF VOWST AND A POSITIVE STOOL TEST FOR C. DIFFICILE), AND (B) HAS NOT RECEIVED MORE THAN ONE TREATMENT COURSE OF VOWST WHICH WAS AT LEAST 12 DAYS AND NOT MORE THAN 8 WEEKS PRIOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FEDRATINIB

Products Affected

INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FENFLURAMINE

Products Affected

FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	DRAVET SYNDROME: INITIAL/RENEWAL: 12 MONTHS. LGS: 12 MONTHS.
Other Criteria	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DRAVET SYNDROME: PATIENT HAS SHOWN CONTINUED CLINICAL BENEFIT (E.G. REDUCTION OF SEIZURES, REDUCED LENGTH OF SEIZURES, SEIZURE CONTROL MAINTAINED).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FENTANYL CITRATE

Products Affected

• fentanyl citrate buccal lozenge on a handle

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE- RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FILGRASTIM-AAFI

Products Affected

NIVESTYM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FILGRASTIM-AYOW

Products Affected

RELEUKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NIVESTYM, WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FILGRASTIM-SNDZ

Products Affected

· ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINERENONE

Products Affected

KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINGOLIMOD

Products Affected

- fingolimod GILENYA ORAL CAPSULE 0.25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FINGOLIMOD LAURYL SULFATE

Products Affected

· TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS (MS): (1) UNABLE TO SWALLOW FINGOLIMOD CAPSULES, AND (2) TRIAL OF OR CONTRAINDICATION TO FINGOLIMOD CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FREMANEZUMAB-VFRM

Products Affected

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

FUTIBATINIB

Products Affected

• LYTGOBI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GALCANEZUMAB-GNLM

Products Affected

EMGALITY PEN

3)

 EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.
Other Criteria	INITIAL: MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: MIGRAINE PREVENTION: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. EPISODIC CLUSTER HEADACHE: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GANAXOLONE

Products Affected

ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GEFITINIB

Products Affected

• gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GILTERITINIB

Products Affected

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLASDEGIB

Products Affected

• DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLATIRAMER

Products Affected

- COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML
- glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml
- glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GLYCEROL PHENYLBUTYRATE

Products Affected

RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	UREA CYCLE DISORDER (UCD): INITIAL: DIAGNOSIS IS CONFIRMED BY ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	UCD: INITIAL: TRIAL OF OR CONTRAINDICATION TO SODIUM PHENYLBUTYRATE. RENEWAL: PATIENT HAS CLINICAL BENEFIT FROM BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GOSERELIN

Products Affected

· ZOLADEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	
Prescriber Restrictions	ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS.
Other Criteria	ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

GUSELKUMAB

Products Affected

TREMFYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

Products Affected

• morphine concentrate oral solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.
Other Criteria	1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - PROMETHAZINE

Products Affected

• promethazine injection solution

mg

- promethazine oral
- promethazine rectal
- promethegan rectal suppository 12.5 mg, 25

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRURITUS/URTICARIA/SEASONAL/PERENNIAL ALLERGY: TRIAL OF OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE OR PRESCRIBER ACKNOWLEDGEMENT OR AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. NAUSEA AND VOMITING: PRESCRIBER ACKNOWLEDGEMENT OR AWARENESS THAT THE DRUG IS CONSIDERED HIGH- RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVES NOR REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - SCOPOLAMINE

Products Affected

• scopolamine base

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - BARBITURATE COMBINATIONS

- ascomp with codeine
- butalbital-acetaminophen-caff oral tablet
- butalbital-aspirin-caffeine
- codeine-butalbital-asa-caff

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - DIPYRIDAMOLE

Products Affected

• dipyridamole oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - DISOPYRAMIDE

Products Affected

• disopyramide phosphate oral capsule

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY -**ENDOCRINE - ESTROGEN**

- amabelz
- dotti
- **DUAVEE**
- estradiol oral
- estradiol transdermal patch semiweekly
- estradiol transdermal patch weekly
- estradiol-norethindrone acet oral tablet 0.5- PREMPHASE 0.1 mg
- fyavolv

- jinteli
- lyllana
- mimvey
- norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg
- PREMARIN ORAL
- PREMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. ALL OTHER FDA APPROVED INDICATIONS NOT PREVIOUSLY MENTIONED IN THIS SECTION, SUCH AS PALLIATIVE TREATMENT, AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - ENDOCRINE - SULFONYLUREAS

- glyburide
- glyburide micronized
- glyburide-metformin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF GLIMEPIRIDE, GLIPIZIDE, OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - KETOROLAC

Products Affected

ketorolac oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	30 DAYS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY - SKELETAL MUSCLE RELAXANTS

- chlorzoxazone oral tablet 500 mg
- cyclobenzaprine oral tablet 10 mg, 5 mg
- methocarbamol oral tablet 500 mg, 750 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED A HIGH RISK MEDICATION FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY ANTICHOLINERGICS -CYPROHEPTADINE_CARBINOXAMINE

Products Affected

• cyproheptadine oral syrup

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-ANTICHOLINERGICS- DIPHENHYDRAMINE ELIXIR

Products Affected

• diphenhydramine hcl oral elixir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ANTIHISTAMINIC CONDITIONS (PRURITUS OR URTICARIA): TRIAL OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE. INSOMNIA: TRIAL OF SILENOR AND BELSOMRA. MOTION SICKNESS AND ANTIPARKINSONISM: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS AND ANAPHYLACTIC REACTIONS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-DIPHENOXYLATE-ATROPINE

Products Affected

• diphenoxylate-atropine oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-INDOMETHACIN

- indomethacin oral capsule 25 mg, 50 mg
- indomethacin oral capsule, extended release

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-MEGESTROL

- megestrol oral suspension 400 mg/10 ml (40 mg/ml)
- megestrol oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK DRUGS IN THE ELDERLY-PAROXETINE

- paroxetine hcl oral suspension
- paroxetine hcl oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HIGH RISK MEDICATIONS IN THE ELDERLY-PHENOBARBITAL

Products Affected

• phenobarbital

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	FOR TREATMENT OF EPILEPSY/SEIZURES IN PATIENTS WHO ARE NEWLY PRESCRIBED PHENOBARBITAL: PATIENT HAS NOT RESPONDED TO AT LEAST ONE ANTICONVULSANT OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

HYDROXYUREA

Products Affected

• SIKLOS ORAL TABLET 100 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IBRUTINIB

Products Affected

• IMBRUVICA ORAL CAPSULE 140 MG, 70 MG

280 MG, 420 MG

- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG,

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IBUPROFEN-FAMOTIDINE

Products Affected

• ibuprofen-famotidine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE OF THE FOLLOWING GENERIC, FEDERAL LEGEND HISTAMINE H2-RECEPTOR ANTAGONISTS: FAMOTIDINE, CIMETIDINE, OR NIZATIDINE, AND TRIAL OF GENERIC, FEDERAL LEGEND IBUPROFEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ICATIBANT

- icatibant
- sajazir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IDELALISIB

Products Affected

• ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IMATINIB

Products Affected

• imatinib oral tablet 100 mg, 400 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
Other Criteria	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFIGRATINIB

Products Affected

• TRUSELTIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INFLIXIMAB

Products Affected

• infliximab

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON ALFA-2B

Products Affected

• INTRON A INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HEPATITIS C: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G. HEPATOLOGIST).
Coverage Duration	6 MONTHS.
Other Criteria	HEPATITIS C: 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) TRIAL OF OR CONTRAINDICATION TO PEGINTERFERON ALFA-2A OR PEGINTERFERON ALFA-2B, 3) USED IN COMBINATION WITH RIBAVIRIN UNLESS CONTRAINDICATED AND 4) LIMITED TO TOTAL OF 24 MONTHS OF TREATMENT. FOLLICULAR LYMPHOMA: LIMITED TO TOTAL OF 18 MONTHS OF TREATMENT. ALL OTHER INDICATIONS: LIMITED TO TOTAL OF 1 YEAR OF TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON FOR MS-AVONEX

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON FOR MS-BETASERON

Products Affected

• BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON FOR MS-PLEGRIDY

Products Affected

 PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML

• PLEGRIDY SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INTERFERON GAMMA-1B

Products Affected

ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IPILIMUMAB

Products Affected

YERVOY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO
Other Criteria	RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IVACAFTOR

Products Affected

KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IVOSIDENIB

Products Affected

• TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IXAZOMIB

Products Affected

NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

IXEKIZUMAB

- TALTZ AUTOINJECTOR
- TALTZ SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA, COSENTYX, XELJANZ, RINVOQ. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, RINVOQ. RENEWAL: PSO, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LANADELUMAB

Products Affected

• TAKHZYRO SUBCUTANEOUS SOLUTION

(150 MG/ML)

 TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LANREOTIDE

Products Affected

• lanreotide

ML

 SOMATULINE DEPOT SUBCUTANEOUS SYRINGE 120 MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP- NETS, CARCINOID SYNDROME: 12 MOS.
Other Criteria	ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LAPATINIB

Products Affected

• lapatinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LAROTRECTINIB

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- · VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	APPROVAL FOR VITRAKVI ORAL SOLUTION: TRIAL OF VITRAKVI CAPSULES OR PATIENT IS UNABLE TO TAKE CAPSULE FORMULATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEDIPASVIR-SOFOSBUVIR

- HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANAVIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. REQUESTS FOR HARVONI 45MG-200MG PELLETS: PATIENT IS UNABLE TO SWALLOW TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LENALIDOMIDE

Products Affected

lenalidomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LENVATINIB

Products Affected

LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LETERMOVIR

- PREVYMIS INTRAVENOUS SOLUTION 240 MG/12 ML, 480 MG/24 ML
- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	HSCT: NOT AT RISK FOR LATE CMV: 4 MOS, AT RISK FOR LATE CMV: 7 MOS. KIDNEY TRANSPLANT: 7 MOS.
Other Criteria	HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION (A) BEYOND 100 DAYS POST TRANSPLANT IF NOT AT RISK FOR LATE CYTOMEGALOVIRUS (CMV) INFECTION AND DISEASE, OR (B) BEYOND 200 DAYS POST TRANSPLANT IF AT RISK FOR LATE CMV INFECTION AND DISEASE. KIDNEY TRANSPLANT: 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 7 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 200 DAYS POST TRANSPLANT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE

Products Affected

• leuprolide subcutaneous kit

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	INITIAL: CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	PROSTATE CANCER: 12 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) PATIENT HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE DEPOT

Products Affected

• leuprolide (3 month)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE-ELIGARD

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LEUPROLIDE-LUPRON DEPOT

- · LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.
Other Criteria	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

LEVODOPA

Products Affected

• INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: 1) NOT CURRENTLY TAKING MORE THAN 1600MG OF LEVODOPA PER DAY, AND 2) PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF INBRIJA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

L-GLUTAMINE

Products Affected

ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME.
Other Criteria	SCD: INITIAL: PATIENTS 18 YEARS OR OLDER: ONE OF THE FOLLOWING: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. PATIENTS 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: HAS MAINTAINED OR EXPERIENCED REDUCTION IN ACUTE COMPLICATIONS OF SICKLE CELL DISEASE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE

- lidocaine hcl mucous membrane solution 4 % (40 mg/ml)
- lidocaine topical adhesive patch, medicated 5
- lidocaine topical ointment
- · ZTLIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LIDOCAINE PRILOCAINE

Products Affected

• lidocaine-prilocaine topical cream

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

LOMITAPIDE

Products Affected

 JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 40 MG, 5 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST.
Coverage Duration	12 MONTHS

PA Criteria	Criteria Details
Other Criteria	1) DIAGNOSIS DETERMINED BY A) DEFINITE SIMON BROOME DIAGNOSTIC CRITERIA, OR B) DUTCH LIPID NETWORK CRITERIA SCORE OF 8 OR GREATER, OR C) CLINICAL DIAGNOSIS BASED ON A HISTORY OF AN UNTREATED LDL-C CONCENTRATION GREATER THAN 500 MG/DL TOGETHER WITH EITHER XANTHOMA BEFORE 10 YEARS OF AGE, OR EVIDENCE OF HEFH IN BOTH PARENTS. 2) LDL-C LEVEL GREATER THAN OR EQUAL TO 70MG/DL WHILE ON MAXIMAL DRUG TREATMENT. 3) TRIAL OF EVOLOCUMAB UNLESS THE PATIENT HAS NON-FUNCTIONING LDL RECEPTORS. 4) MEETS ONE OF THE FOLLOWING: A) TAKING A HIGH-INTENSITY STATIN (I.E., ATORVASTATIN 40-80MG DAILY, ROSUVASTATIN 20-40MG DAILY) FOR A DURATION OF AT LEAST 8 WEEKS, B) TAKING A MAXIMALLY TOLERATED DOSE OF ANY STATIN FOR A DURATION OF AT LEAST 8 WEEKS GIVEN THAT THE PATIENT CANNOT TOLERATE A HIGH-INTENSITY STATIN, C) ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (E.G., ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTIONS), D) STATIN INTOLERANCE, OR E) TRIAL OF ROSUVASTATIN, ATORVASTATIN, OR STATIN THERAPY AT ANY DOSE AND HAS EXPERIENCED SKELETAL-
Indications	MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY). All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LONCASTUXIMAB TESIRINE-LPYL

Products Affected

· ZYNLONTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LORLATINIB

Products Affected

 LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LUMACAFTOR-IVACAFTOR

- ORKAMBI ORAL GRANULES IN PACKET
- · ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

LUMASIRAN

Products Affected

OXLUMO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MACITENTAN

Products Affected

• OPSUMIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR 2) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MARGETUXIMAB-CMKB

Products Affected

MARGENZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MEPOLIZUMAB

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: ASTHMA: 4 MO. NASAL POLYPS: 6 MO. OTHERS: 12 MO. RENEWAL: NASAL POLYPS, ASTHMA: 12 MO.

PA Criteria	Criteria Details
Other Criteria	INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. NASAL POLYPS: PREVIOUS 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. RENEWAL: ASTHMA: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEVI FROM
Indications	PRETREATMENT BASELINE. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE. All EDA-approved Indications
Off Label Uses	All FDA-approved Indications.

PA Criteria	Criteria Details
Part B Prerequisite	No

MIDOSTAURIN

Products Affected

RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIFEPRISTONE

Products Affected

KORLYM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM).
Age Restrictions	
Prescriber Restrictions	CS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO KORLYM, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIGALASTAT

Products Affected

GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	FABRY DISEASE: INITIAL: PATIENT IS SYMPTOMATIC OR HAS EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS.
Age Restrictions	
Prescriber Restrictions	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MOS. RENEWAL: 12 MOS.
Other Criteria	FABRY DISEASE: INITIAL: NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME), RENEWAL: 1) PATIENT HAS DEMONSTRATED IMPROVEMENT OR STABILIZATION, AND 2) NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MIGLUSTAT

- miglustat
- yargesa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MILTEFOSINE

Products Affected

• IMPAVIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MOBOCERTINIB

Products Affected

EXKIVITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MOMELOTINIB

Products Affected

OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

MOSUNETUZUMAB-AXGB

Products Affected

LUNSUMIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS.
Other Criteria	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NAFARELIN

Products Affected

• SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ENDOMETRIOSIS: 6 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. CPP: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: CPP: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NARCOLEPSY AGENTS

Products Affected

- armodafinil
- modafinil oral tablet 100 mg, 200 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

NATALIZUMAB

Products Affected

TYSABRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	MULTIPLE SCLEROSIS (MS): 12 MOS. CD: INITIAL: 6 MOS, RENEWAL: 12 MOS.
Other Criteria	INITIAL: MS: TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF MS. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI, RINVOQ. RENEWAL: CD: ONE OF THE FOLLOWING: 1) RECEIVED AT LEAST 12 MONTHS OF THERAPY WITH TYSABRI AND HAS NOT REQUIRED MORE THAN 3 MONTHS OF CORTICOSTEROID USE WITHIN THE PAST 12 MONTHS TO CONTROL THEIR CROHNS DISEASE WHILE ON TYSABRI, OR 2) HAS ONLY RECEIVED 6 MONTHS OF THERAPY WITH TYSABRI AND HAS TAPERED OFF CORTICOSTEROIDS DURING THE FIRST 24 WEEKS OF TYSABRI THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

NAXITAMAB-GQGK

Products Affected

DANYELZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NERATINIB MALEATE

Products Affected

NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NILOTINIB

Products Affected

 TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND TASIGNA IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NINTEDANIB

Products Affected

• OFEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) BASELINE FORCED VITAL CAPACITY (FVC) AT LEAST 50% OF PREDICTED VALUE. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 45% OF PREDICTED VALUE.
Age Restrictions	
Prescriber Restrictions	INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG-INDUCED LUNG TOXICITY, RECURRENT ASPIRATION), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ACTEMRA SUBQ. PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENED/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIRAPARIB

Products Affected

- · ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIRAPARIB-ABIRATERONE

Products Affected

AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NITISINONE

Products Affected

- nitisinone
- · ORFADIN ORAL CAPSULE 20 MG
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.
Age Restrictions	
Prescriber Restrictions	HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIVOLUMAB

Products Affected

• OPDIVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

NIVOLUMAB-RELATLIMAB-RMBW

Products Affected

OPDUALAG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OBETICHOLIC ACID

Products Affected

· OCALIVA

PA Criteria	Criteria Details
Exclusion	PRIMARY BILIARY CHOLANGITIS (PBC):
Criteria	INITIAL/RENEWAL: COMPLETE BILIARY OBSTRUCTION.
Required Medical	PBC: INITIAL: DIAGNOSIS CONFIRMED BY TWO OF THE
Information	FOLLOWING: 1) ALKALINE PHOSPHATASE LEVEL OF AT
	LEAST 1.5 TIMES THE UPPER LIMIT OF NORMAL, 2)
	PRESENCE OF ANTIMITOCHONDRIAL ANTIBODIES AT A
	TITER OF 1:40 OR HIGHER, OR 3) HISTOLOGIC EVIDENCE
	OF NON-SUPPURATIVE DESTRUCTIVE CHOLANGITIS AND
	DESTRUCTION OF INTERLOBULAR BILE DUCTS.
Age Restrictions	
Prescriber	PBC: INITIAL: PRESCRIBED BY OR IN CONSULTATION
Restrictions	WITH A GASTROENTEROLOGIST OR HEPATOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	PBC: INITIAL: USED IN COMBINATION WITH
Other Criteria	URSODEOXYCHOLIC ACID IN A PATIENT WITH AN
	INADEQUATE RESPONSE TO URSODEOXYCHOLIC ACID,
	OR AS MONOTHERAPY IN A PATIENT WHO IS UNABLE TO
	TOLERATE URSODEOXYCHOLIC ACID. RENEWAL:
	CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OCRELIZUMAB

Products Affected

• OCREVUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OFATUMUMAB-SQ

Products Affected

KESIMPTA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OLANZAPINE/SAMIDORPHAN

Products Affected

LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SCHIZOPHRENIA/BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST
Coverage Duration	12 MONTHS
Other Criteria	SCHIZOPHRENIA: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO LATUDA OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OLAPARIB

Products Affected

· LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: 1) MEDICATION WILL BE USED AS MONOTHERAPY, AND 2) PATIENT HAS COMPLETED TWO OR MORE LINES OF PLATINUM-BASED CHEMOTHERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OLUTASIDENIB

Products Affected

REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OMACETAXINE

Products Affected

• SYNRIBO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

OMALIZUMAB

Products Affected

XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL GREATER THAN OR EQUAL TO 30 IU/ML.
Age Restrictions	
Prescriber Restrictions	INITIAL AND RENEWAL: CHRONIC IDIOPATHIC URTICARIA (CIU): PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE, DERMATOLOGY OR IMMUNOLOGY. INITIAL: NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: ASTHMA: 12 MO. CIU, NASAL POLYPS: 6 MO. RENEWAL: ASTHMA, NASAL POLYPS: 12 MO. CIU: 6 MO.

PA Criteria	Criteria Details
Other Criteria	INITIAL: CIU: TRIAL OF OR CONTRAINDICATION TO A MAXIMALLY TOLERATED DOSE OF AN HI ANTI-HISTAMINE AND STILL EXPERIENCES HIVES ON MOST DAYS OF THE WEEK. NASAL POLYPS: 1) PREVIOUS 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) NOT CONCURRENTLY RECEIVING DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: CIU: DIAGNOSIS OF CIU. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE. ASTHMA: 1) NOT CONCURRENTLY RECEIVING DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: A) REDUCTION IN ASTHMA
	EXACERBATIONS FROM BASELINE, B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

OSIMERTINIB

Products Affected

TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EGFR EXON 19 DELETIONS OR EXON 21 L858R MUTATIONS NON-SMALL CELL LUNG CANCER (NSCLC) AND METASTATIC NSCLC WITH EGFR T790M MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PACRITINIB

Products Affected

· VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PALBOCICLIB

Products Affected

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PANOBINOSTAT

Products Affected

FARYDAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	MULTIPLE MYELOMA: RENEWAL: TOLERATED THE FIRST 8 CYCLES OF THERAPY WITHOUT UNRESOLVED SEVERE OR MEDICALLY SIGNIFICANT TOXICITY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PARATHYROID HORMONE

Products Affected

NATPARA

PA Criteria	Criteria Details
Exclusion	
Criteria	
Required Medical Information	
Age Restrictions	
Prescriber	HYPOCALCEMIA SECONDARY TO
Restrictions	HYPOPARATHYROIDISM: PRESCRIBED BY OR IN
	CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage	12 MONTHS
Duration	
Other Criteria	HYPOCALCEMIA SECONDARY TO
	HYPOPARATHYROIDISM: 1) TRIAL OF OR
	CONTRAINDICATION TO CALCITRIOL, 2)
	HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM
	SENSING RECEPTOR (CSR) MUTATION, AND 3)
	HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE
	POST-SURGICAL HYPOPARATHYROIDISM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PASIREOTIDE DIASPARTATE

Products Affected

SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PAZOPANIB

Products Affected

- pazopanib VOTRIENT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PDE5 INHIBITORS FOR PULMONARY ARTERIAL HYPERTENSION

Products Affected

- alyq
- sildenafil (pulm.hypertension) oral tablet
- tadalafil (pulm. hypertension)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION, 2) PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS. THESE CRITERIA DO NOT APPLY TO SILDENAFIL FOR AGES 1 TO 17 YEARS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. THIS CRITERIA DOES NOT APPLY TO SILDENAFIL FOR AGES 1 TO 17 YEARS.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	PAH: INITIAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS, AND 2) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER. RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS, AND 2) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS. ALL OF THE PRECEDING CRITERIA DOES NOT APPLY TO SILDENAFIL FOR AGES 1 TO 17 YEARS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM

Products Affected

NEULASTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM - APGF

Products Affected

NYVEPRIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM - CBQV

Products Affected

- UDENYCA
- UDENYCA AUTOINJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM - JMDB

Products Affected

FULPHILA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGFILGRASTIM-NEULASTA ONPRO

Products Affected

NEULASTA ONPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGVALIASE-PQPZ

Products Affected

PALYNZIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	PHENYLKETONURIA (PKU): INITIAL: NOT ON CONCURRENT TREATMENT WITH KUVAN. RENEWAL: 1) PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NOT ON CONCURRENT TREATMENT WITH KUVAN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEGVISOMANT

Products Affected

SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEMBROLIZUMAB

Products Affected

KEYTRUDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEMIGATINIB

Products Affected

PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PENICILLAMINE

Products Affected

• penicillamine oral tablet

PA Criteria	Criteria Details
Exclusion	
Criteria	
Required Medical	INITIAL: WILSONS DISEASE: CONFIRMED BY ONE OF THE
Information	FOLLOWING: 1) PLASMA COPPER-PROTEIN
	CERULOPLASMIN IS LESS THAN 20MG/DL, 2) LIVER
	BIOPSY POSITIVE FOR AN ABNORMALLY HIGH
	CONCENTRATION OF COPPER (GREATER THAN 250MCG/G
	DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER
	RINGS, OR 3) CONFIRMATION BY GENETIC TESTING FOR
	ATP7B MUTATIONS. CYSTINURIA: PATIENT HAS
	NEPHROLITHIASIS AND ONE OR MORE OF THE
	FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE
	OF CYSTEINE, 2) IDENTIFICATION OF PATHOGNOMONIC
	HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3)
	POSITIVE FAMILY HISTORY OF CYSTINURIA WITH
	POSITIVE CYANIDE-NITROPRUSSIDE SCREEN.
Age Restrictions	
Prescriber	WILSONS DISEASE: PRESCRIBED BY OR IN
Restrictions	CONSULTATION WITH A HEPATOLOGIST OR
	GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY
	OR IN CONSULTATION WITH A NEPHROLOGIST.
	RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN
	CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.

PA Criteria	Criteria Details
Other Criteria	INITIAL: WILSONS DISEASE: 1) KNOWN FAMILY HISTORY OF WILSONS DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSONS DISEASE, AND 2) REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRE A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN). CYSTINURIA: REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRES A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN) AND A FORMULARY VERSION OF TIOPRONIN (THIOLA)/THIOLA EC. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED, AND 3) REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRES A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN). RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PEXIDARTINIB

Products Affected

TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PIMAVANSERIN

Products Affected

NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER
Prescriber Restrictions	PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST).
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PIRFENIDONE

Products Affected

- pirfenidone oral capsule
- pirfenidone oral tablet 267 mg, 534 mg, 801 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50% AT BASELINE.
Age Restrictions	IPF: INITIAL: 18 YEARS OR OLDER.
Prescriber Restrictions	IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

PIRTOBRUTINIB

Products Affected

• JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

POMALIDOMIDE

Products Affected

POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PONATINIB

Products Affected

• ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

POSACONAZOLE

Products Affected

• posaconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OROPHARYNGEAL CANDIDIASIS (OPC): 3 MONTHS. PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS.
Other Criteria	POSACONAZOLE SUSPENSION ONLY: 1) OPC: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE OR ITRACONAZOLE. 2) PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. POSACONZOLE TABLETS ONLY: TREATMENT OF INVASIVE ASPERGILLOSIS: NO EXTRA CRITERIA REQUIRED. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

POSACONAZOLE-POWDERMIX

Products Affected

• NOXAFIL ORAL SUSP, DELAYED RELEASE FOR RECON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PRALSETINIB

Products Affected

• GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PRAMLINTIDE

Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TYPE I OR TYPE II DIABETES: REQUIRING INSULIN OR CONTINUOUS INSULIN INFUSION (INSULIN PUMP) FOR GLYCEMIC CONTROL
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PYRIMETHAMINE

Products Affected

• pyrimethamine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.
Other Criteria	TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

QUININE

Products Affected

• quinine sulfate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

QUIZARTINIB

Products Affected

VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RAMUCIRUMAB

Products Affected

· CYRAMZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

REGORAFENIB

Products Affected

STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RELUGOLIX

Products Affected

ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RESLIZUMAB

Products Affected

· CINQAIR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	ASTHMA: INITIAL 4 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: FASENRA, NUCALA, DUPIXENT, AND 4) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEVI FROM PRETREATMENT BASELINE. THIS DRUG ALSO REQUIRES
	PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

RETIFANLIMAB-DLWR

Products Affected

ZYNYZ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIBOCICLIB

Products Affected

- KISQALI FEMARA CO-PACK ORAL KISQALI ORAL TABLET 200 TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG
 - MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO VERZENIO OR IBRANCE WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIFAXIMIN

Products Affected

• XIFAXAN ORAL TABLET 200 MG, 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	TRAVELERS DIARRHEA/HE: 12 MOS. IBS-D: 8 WKS.
Other Criteria	RIFAXIMIN 550 MG TABLETS: HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIMEGEPANT

Products Affected

· NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: ACUTE MIGRAINE TREATMENT: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: ACUTE MIGRAINE TREATMENT: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

RIOCIGUAT

Products Affected

ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): 1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) WHO GROUP 4: NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: PAH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, OR ANY PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS, AND 2) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RIPRETINIB

Products Affected

· QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RISANKIZUMAB-RZAA

Products Affected

SKYRIZI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

RISDIPLAM

Products Affected

• EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	SPINAL MUSCULAR ATROPHY (SMA): INITIAL: GENE MUTATION ANALYSIS INDICATING MUTATIONS OR DELETIONS OF BOTH ALLELES OF THE SURVIVAL MOTOR NEURON 1 (SMN1) GENE. FOR PRESYMPTOMATIC PATIENTS: UP TO THREE COPIES OF SURVIVAL MOTOR NEURON 2 (SMN2) BASED ON NEWBORN SCREENING.
Age Restrictions	
Prescriber Restrictions	SMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST AT A SMA SPECIALTY CENTER.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	SMA: INITIAL: FOR SYMPTOMATIC PATIENTS: (1) BASELINE MOTOR FUNCTION ASSESSMENT BY A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST, AND (2) IF PATIENT RECEIVED GENE THERAPY, THE PATIENT HAD LESS THAN EXPECTED CLINICAL BENEFIT. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN: (1) MOTOR FUNCTION ASSESSMENTS COMPARED TO BASELINE, OR (2) OTHER MUSCLE FUNCTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB AND HYALURONIDASE HUMAN-SQ

Products Affected

RITUXAN HYCELA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PATIENT HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB-ABBS

Products Affected

TRUXIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB-ARRX

Products Affected

RIABNI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, WG, MPA: 12 MO. CLL: 6 MO.
Other Criteria	RA: INITIAL: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RITUXIMAB-PVVR

Products Affected

RUXIENCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
Coverage Duration	NHL, GPA, MPA: 12 MONTHS. CLL: 6 MONTHS. RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO PREFERRED AGENTS: HUMIRA, ENBREL, RINVOQ, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ROPEGINTERFERON ALFA-2B-NJFT

Products Affected

BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RUCAPARIB

Products Affected

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

RUXOLITINIB

Products Affected

JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS.
Other Criteria	MYELOFIBROSIS: RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SAPROPTERIN

Products Affected

- javygtor oral tablet, soluble
- sapropterin oral tablet, soluble

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 1 MONTH, RENEWAL 12 MONTHS.
Other Criteria	HYPERPHENYLALANINEMIA (HPA): INITIAL: NOT CONCURRENTLY USING PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NOT CONCURRENTLY USING PALYNZIQ.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SARILUMAB

Products Affected

KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYMYALGIA RHEUMATICA (PMR): 12 MONTHS.
Other Criteria	RA: INITIAL: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SECUKINUMAB

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML

• COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI-RHEUMATIC DRUG). AS, NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. RENEWAL: PSO, PSA, AS, NR-AXSPA, ERA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELEXIPAG

Products Affected

- UPTRAVI INTRAVENOUS
- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS, DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	PAH: INITIAL: WHO FC II-III SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS, EACH FROM A DIFFERENT DRUG CLASS: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. WHO FC III SYMPTOMS AND EVIDENCE OF RAPID PROGRESSION OR POOR PROGNOSIS, WHO FC IV SYMPTOMS: NO STEP. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST, OR 2) REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELINEXOR

Products Affected

XPOVIO ORAL TABLET 100
 MG/WEEK (50 MG X 2), 40 MG/WEEK
 (40 MG X 1), 40MG TWICE WEEK (40
 MG X 2), 60 MG/WEEK (60 MG X 1),
 60MG TWICE WEEK (120 MG/WEEK),
 80 MG/WEEK (40 MG X 2), 80MG
 TWICE WEEK (160 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELPERCATINIB

Products Affected

• RETEVMO ORAL CAPSULE 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SELUMETINIB

Products Affected

 KOSELUGO ORAL CAPSULE 10 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SIPONIMOD

Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1
 MG, 2 MG
 MAYZENT STARTER(FOR 2MG MAINT)
- MAYZENT STARTER(FOR 1MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS: RENEWAL: 1) DEMONSTRATION OF CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE AND 2) DOES NOT HAVE LYMPHOPENIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SIROLIMUS PROTEIN-BOUND

Products Affected

FYARRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SODIUM OXYBATE

Products Affected

- sodium oxybate XYREM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: 1) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT, 2) FOR PATIENTS 18 YEARS OR OLDER: TRIAL OF OR CONTRAINDICATION TO THE FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, PITOLISANT OR SOLRIAMFETOL AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) FOR PATIENTS 7 TO 17 YEARS OF AGE: TRIAL OF OR CONTRAINDICATION TO ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT. RENEWAL (ALL INDICATIONS): 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

SOFOSBUVIR/VELPATASVIR

Products Affected

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANAVIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

Products Affected

VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

SOLRIAMFETOL

Products Affected

• SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: EDS IN NARCOLEPSY: TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL, AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. EDS IN OBSTRUCTIVE SLEEP APNEA (OSA): TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL. RENEWAL: EDS IN NARCOLEPSY OR OSA: SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOMATROPIN - NORDITROPIN

Products Affected

NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES.
Required Medical Information	INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: ADULT GHD: GROWTH HORMONE DEFICIENCY ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASES, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, TRAUMA, OR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GROWTH HORMONE DEFICIENCY. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR PATIENT HAS NOT COMPLETED PREPUBERTALGROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOMATROPIN - SEROSTIM

Products Affected

• SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI- AGING PURPOSES
Required Medical Information	INITIAL: HIV/WASTING: MEETS ONE OF THE FOLLOWING CRITERIA FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BODY CELL MASS (BCM) LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED.
Age Restrictions	
Prescriber Restrictions	HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	INITIAL/RENEWAL: 3 MONTHS.
Other Criteria	HIV/WASTING: INITIAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

SONIDEGIB

Products Affected

· ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC): BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SORAFENIB

Products Affected

• sorafenib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SOTORASIB

Products Affected

• LUMAKRAS ORAL TABLET 120 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

STIRIPENTOL

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

SUNITINIB

Products Affected

• sunitinib malate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TALAZOPARIB

Products Affected

TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: PATIENT HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING. PATIENTS WITH HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER MUST HAVE ADDITIONAL PRIOR TREATMENT WITH ENDOCRINE THERAPY OR BE CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TALIMOGENE

Products Affected

• IMLYGIC INJECTION SUSPENSION 10EXP6 (1 MILLION) PFU/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE MELANOMA: 1) IMLYGIC TO BE INJECTED INTO CUTANEOUS, SUBCUTANEOUS, AND/OR NODAL LESIONS THAT ARE VISIBLE, PALPABLE, OR DETECTABLE BY ULTRASOUND GUIDANCE, 2) NOT CURRENTLY RECEIVING IMMUNOSUPPRESSIVE THERAPY, 3) NO HISTORY OF PRIMARY OR ACQUIRED IMMUNODEFICIENT STATES, LEUKEMIA, LYMPHOMA, OR AIDS, AND 4) NO CONCURRENT USE WITH PEMBROLIZUMAB, NIVOLUMAB, IPILIMUMAB, DABRAFENIB, TRAMETINIB, VEMURAFENIB, INTERLEUKIN-2, INTERFERON, DACARBAZINE, TEMOZOLOMIDE, PACLITAXEL, CARBOPLATIN, IMATINIB, MELPHALAN, IMIQUIMOD, OR RADIATION THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TALQUETAMAB-TGVS

Products Affected

TALVEY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TASIMELTEON

Products Affected

tasimelteon

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	LIFETIME
Other Criteria	NON-24 HOUR SLEEP-WAKE DISORDER: PATIENT IS LIGHT-INSENSITIVE OR HAS TOTAL BLINDNESS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TAZEMETOSTAT

Products Affected

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TBO-FILGRASTIM

Products Affected

• GRANIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON-MYELOID MALIGNANCIES: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON-MYELOID MALIGNANCIES: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEBENTAFUSP-TEBN

Products Affected

KIMMTRAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TECLISTAMAB-CQYV

Products Affected

TECVAYLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEDUGLUTIDE

Products Affected

• GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SHORT BOWEL SYNDROME (SBS): INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	SBS: INITIAL: PATIENT IS DEPENDENT ON INTRAVENOUS PARENTERAL NUTRITION DEFINED AS REQUIRING PARENTERAL NUTRITION AT LEAST THREE TIMES PER WEEK. RENEWAL: ACHIEVED OR MAINTAINED A DECREASED NEED FOR PARENTERAL SUPPORT COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TELOTRISTAT

Products Affected

XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEPOTINIB

Products Affected

TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TERIFLUNOMIDE

Products Affected

• teriflunomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TESAMORELIN

Products Affected

• EGRIFTA SV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TESTOSTERONE

- testosterone cypionate
- testosterone enanthate
- testosterone transdermal gel in metereddose pump 12.5 mg/ 1.25 gram (1%), 20.25 mg/1.25 gram (1.62%)
- testosterone transdermal gel in packet 1 % (25 mg/2.5gram), 1 % (50 mg/5 gram)
- testosterone transdermal solution in metered pump wlapp
- XYOSTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: MALE HYPOGONADISM: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	MALE HYPOGONADISM: INITIAL/RENEWAL:12 MO. ALL OTHER INDICATIONS: LIFETIME OF MEMBERSHIP IN PLAN.
Other Criteria	RENEWAL: MALE HYPOGONADISM: IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TETRABENAZINE

Products Affected

• tetrabenazine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TEZACAFTOR/IVACAFTOR

Products Affected

SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

THALIDOMIDE

Products Affected

• THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TISOTUMAB VEDOTIN-TFTV

Products Affected

TIVDAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TIVOZANIB

Products Affected

FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TOCILIZUMAB IV

Products Affected

ACTEMRA

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
Coverage Duration	INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

PA Criteria	Criteria Details
Off Label Uses	
Part B Prerequisite	No

TOCILIZUMAB SQ

- ACTEMRA
- ACTEMRA ACTPEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TOFACITINIB

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA, PCJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, PCJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TOPICAL TRETINOIN

- ALTRENO
- tretinoin

PA Criteria	Criteria Details
Exclusion Criteria	COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRAMETINIB

Products Affected

 MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRAMETINIB SOLUTION

Products Affected

• MEKINIST ORAL RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNABLE TO SWALLOW MEKINIST TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB HYALURONIDASE

Products Affected

HERCEPTIN HYLECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-DKST

Products Affected

• OGIVRI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-DTTB

Products Affected

ONTRUZANT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-PKRB

Products Affected

HERZUMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRASTUZUMAB-QYYP

Products Affected

TRAZIMERA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TREMELIMUMAB-ACTL

Products Affected

• IMJUDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS.
Other Criteria	UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRIENTINE

Products Affected

• trientine oral capsule 250 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	WILSONS DISEASE: INITIAL: KNOWN FAMILY HISTORY OF WILSONS DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSONS DISEASE. CONFIRMATION OF ONE OF THE FOLLOWING: 1) PLASMA COPPER-PROTEIN CERULOPLASMIN LESS THAN 20 MG/DL, 2) LIVER BIOPSY POSITIVE FOR AN ABNORMALLY HIGH CONCENTRATION OF COPPER (GREATER THAN 250 MCG/G DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER RINGS, OR 3) CONFIRMATION BY GENETIC TESTING FOR ATP7B MUTATIONS.
Age Restrictions	
Prescriber Restrictions	WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
Other Criteria	WILSONS DISEASE: INITIAL: TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE (DEPEN). RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRIFLURIDINE/TIPIRACIL

Products Affected

 LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TRIPTORELIN-TRELSTAR

Products Affected

 TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

TUCATINIB

Products Affected

• TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

UBROGEPANT

Products Affected

• UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

UPADACITINIB

Products Affected

· RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: RA, AS, NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. UC, CD: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE- MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. ATOPIC DERMATITIS: 1) ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING: TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. ATOPIC DERMATITIS: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER
Indications	SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS. All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

USTEKINUMAB

Products Affected

• STELARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

USTEKINUMAB IV

Products Affected

• STELARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	2 MONTHS
Other Criteria	CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VANDETANIB

Products Affected

• CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VEMURAFENIB

Products Affected

• ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VENETOCLAX

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VIGABATRIN

Products Affected

- vigabatrin
- vigadrone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	CPS: 1) TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS AND 2) BENEFITS OUTWEIGH THE POTENTIAL FOR VISION LOSS. INFANTILE SPASMS: BENEFITS OUTWEIGH THE POTENTIAL FOR VISION LOSS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VISMODEGIB

Products Affected

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

VORICONAZOLE SUSPENSION

Products Affected

• voriconazole oral suspension for reconstitution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CANDIDA INFECTIONS: 3 MOS. ALL OTHER INDICATIONS: 6 MOS.
Other Criteria	CANDIDA INFECTIONS: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE. ALL INDICATIONS: INABILITY TO SWALLOW TABLETS OR AN INDICATION FOR ESOPHAGEAL CANDIDIASIS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

ZANUBRUTINIB

Products Affected

• BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

INDEX

abiraterone6	BETASERON SUBCUTANEOUS KIT
ACTEMRA346, 348	
ACTEMRA ACTPEN 348	<i>bexarotene</i> 45
ACTHAR68	bortezomib injection recon soln 1 mg, 2.5
ACTIMMUNE174	<i>mg</i> 47
ADEMPAS288	BORTEZOMIB INTRAVENOUS
AJOVY AUTOINJECTOR130	RECON SOLN
AJOVY SYRINGE130	BOSULIF ORAL TABLET 100 MG,
AKEEGA229	400 MG, 500 MG 50
ALECENSA13	BRAFTOVI ORAL CAPSULE 75 MG 108
ALTRENO352	BRUKINSA
ALUNBRIG ORAL TABLET 180 MG,	butalbital-acetaminophen-caff oral tablet 145
30 MG, 90 MG51	butalbital-aspirin-caffeine145
ALUNBRIG ORAL TABLETS, DOSE	CABLIVI INJECTION KIT57
PACK51	CABOMETYX ORAL TABLET 20
<i>alyq</i>	MG, 40 MG, 60 MG 55
<i>amabelz</i>	CALQUENCE8
<i>ambrisentan</i> 15	CALQUENCE (ACALABRUTINIB
<i>apomorphine</i>	MAL)8
armodafinil220	CAPRELSA ORAL TABLET 100 MG,
ascomp with codeine145	300 MG372
AUSTEDO ORAL TABLET 12 MG, 6	<i>carglumic acid</i> 59
MG, 9 MG85	CAYSTON31
AUSTEDO XR ORAL TABLET	CERDELGA102
EXTENDED RELEASE 24 HR 12	chlorzoxazone oral tablet 500 mg152
MG, 24 MG, 6 MG 85	CIMZIA61
AUSTEDO XR TITRATION	CIMZIA POWDER FOR RECONST61
KT(WK1-4)85	CINQAIR280
AVONEX INTRAMUSCULAR PEN	CINRYZE52
INJECTOR KIT171	codeine-butalbital-asa-caff145
AVONEX INTRAMUSCULAR	colchicine oral tablet67
SYRINGE KIT 171	COMETRIQ ORAL CAPSULE 100
AYVAKIT27	MG/DAY(80 MG X1-20 MG X1), 140
BALVERSA ORAL TABLET 3 MG, 4	MG/DAY(80 MG X1-20 MG X3), 60
MG, 5 MG114	MG/DAY (20 MG X 3/DAY)54
bendamustine intravenous recon soln38	COPAXONE SUBCUTANEOUS
BENDAMUSTINE INTRAVENOUS	SYRINGE 20 MG/ML, 40 MG/ML 137
SOLUTION38	COPIKTRA97
BENDEKA	CORTROPHIN GEL68
BENLYSTA SUBCUTANEOUS35	COSENTYX (2 SYRINGES)303
BESREMI	COSENTYX PEN (2 PENS)303
betaine41	COSENTYX SUBCUTANEOUS
	SYRINGE 75 MG/0.5 ML

COSENTYX UNOREADY PEN 303	EMGALITY SYRINGE	
COTELLIC	SUBCUTANEOUS SYRINGE 120	
cyclobenzaprine oral tablet 10 mg, 5 mg 152	MG/ML, 300 MG/3 ML (100 MG/ML	
cyproheptadine oral syrup153	X 3)	32
CYRAMZA277	ENBREL1	17
CYSTARAN71	ENBREL MINI1	17
<i>dalfampridine</i>	ENBREL SURECLICK1	17
DANYELZA223	ENDARI1	96
DAURISMO ORAL TABLET 100	EPCLUSA ORAL PELLETS IN	
MG, 25 MG136	PACKET 150-37.5 MG, 200-50 MG 3	314
<i>deferasirox</i> 80	EPCLUSA ORAL TABLET3	314
<i>deferiprone</i>	EPIDIOLEX	56
DIACOMIT ORAL CAPSULE 250	EPKINLY1	11
MG, 500 MG325	ERBITUX	63
DIACOMIT ORAL POWDER IN	ERIVEDGE3	376
PACKET 250 MG, 500 MG325	ERLEADA ORAL TABLET 240 MG,	
diclofenac sodium topical gel 3 %87	60 MG	17
diclofenac sodium topical solution in	erlotinib oral tablet 100 mg, 150 mg, 25	
metered-dose pump	<i>mg</i> 1	
dimethyl fumarate oral capsule,delayed	estradiol oral1	
release(drlec) 120 mg, 120 mg (14)- 240	estradiol transdermal patch semiweekly1	
<i>mg</i> (46), 240 <i>mg</i> 89	estradiol transdermal patch weekly1	.48
diphenhydramine hcl oral elixir154	estradiol-norethindrone acet oral tablet	
diphenoxylate-atropine oral tablet156	0.5-0.1 mg1	.48
dipyridamole oral146	everolimus (antineoplastic) oral tablet 10	
disopyramide phosphate oral capsule 147	<i>mg</i> , 2.5 <i>mg</i> , 5 <i>mg</i> , 7.5 <i>mg</i>	.19
DOPTELET (10 TAB PACK)28	everolimus (antineoplastic) oral tablet for	
DOPTELET (15 TAB PACK)28	suspension1	
DOPTELET (30 TAB PACK)28	EVRYSDI2	
<i>dotti</i> 148	EXKIVITY2	
dronabinol92	FARYDAK2	
droxidopa93	FASENRA	
DUAVEE	FASENRA PEN	39
DUPIXENT PEN	fentanyl citrate buccal lozenge on a	
DUPIXENT SYRINGE94	handle1	
EGRIFTA SV339	FERRIPROX (2 TIMES A DAY)	
ELIGARD192	FERRIPROX ORAL SOLUTION	
ELIGARD (3 MONTH)	fingolimod1	
ELIGARD (4 MONTH)	FINTEPLA1	
ELIGARD (6 MONTH) 192	FOTIVDA3	
ELREXFIO 44 MG/1.1 ML VIAL103	FULPHILA2	
ELREXFIO SUBCUTANEOUS	FYARRO3	
SOLUTION 40 MG/ML	<i>fyavolv</i> 1	
EMGALITY PEN132	GALAFOLD2	
	GAMIFANT1	05

GATTEX 30-VIAL	335	IMBRUVICA ORAL CAPSULE 140	
GAVRETO	272	MG, 70 MG	162
gefitinib	134	IMBRUVICA ORAL SUSPENSION	162
GILENYA ORAL CAPSULE 0.25 M	ЛG	IMBRUVICA ORAL TABLET 140	
	128	MG, 280 MG, 420 MG	162
GILOTRIF		IMJUDO	
glatiramer subcutaneous syringe 20		IMLYGIC INJECTION SUSPENSION	1
mg/ml, 40 mg/ml	137	10EXP6 (1 MILLION) PFU/ML	328
glatopa subcutaneous syringe 20 mg/m	l,	IMPAVIDO	.214
40 mg/ml	137	INBRIJA INHALATION CAPSULE,	
glyburide	150	W/INHALATION DEVICE	195
glyburide micronized	150	indomethacin oral capsule 25 mg, 50 mg	157
glyburide-metformin	150	indomethacin oral capsule, extended	
GRANIX	332	release	157
HAEGARDA SUBCUTANEOUS		infliximab	168
RECON SOLN 2,000 UNIT, 3,000		INLYTA ORAL TABLET 1 MG, 5	
UNIT	53	MG	29
HARVONI ORAL PELLETS IN		INQOVI	79
PACKET 33.75-150 MG, 45-200 MG	i 185	INREBIC	121
HARVONI ORAL TABLET	185	INTRON A INJECTION RECON	
HERCEPTIN HYLECTA	355	SOLN	170
HERZUMA	358	JAKAFI	.300
HUMIRA PEN	10	javygtor oral tablet, soluble	301
HUMIRA PEN CROHNS-UC-HS		JAYPIRCA ORAL TABLET 100 MG,	
START	10	50 MG	267
HUMIRA PEN PSOR-UVEITS-AD	OL	JEMPERLI	91
HS	10	jinteli	148
HUMIRA SUBCUTANEOUS		JUXTAPID ORAL CAPSULE 10 MG,	,
SYRINGE KIT 40 MG/0.8 ML	10	20 MG, 30 MG, 40 MG, 5 MG, 60 MG	199
HUMIRA(CF)	10	KALYDECO	.176
HUMIRA(CF) PEDI CROHNS		KERENDIA	.127
STARTER		KESIMPTA PEN	235
HUMIRA(CF) PEN	10	ketorolac oral	151
HUMIRA(CF) PEN CROHNS-UC-	HS.10	KEVZARA	.302
HUMIRA(CF) PEN PEDIATRIC U	C10	KEYTRUDA	259
HUMIRA(CF) PEN PSOR-UV-ADO	OL	KIMMTRAK	.333
HS	10	KISQALI FEMARA CO-PACK ORAI	∟
IBRANCE	245	TABLET 200 MG/DAY(200 MG X 1)-	
ibuprofen-famotidine	163	2.5 MG, 400 MG/DAY(200 MG X 2)-	
icatibant		2.5 MG, 600 MG/DAY(200 MG X 3)-	
ICLUSIG	269	2.5 MG	284
IDHIFA	107	KISQALI ORAL TABLET 200	
IGALMI	86	MG/DAY (200 MG X 1), 400 MG/DAY	7
imatinib oral tablet 100 mg, 400 mg	166	(200 MG X 2), 600 MG/DAY (200 MG	
		X 3)	284

KORLYM211	MAYZENT STARTER(FOR 1MG	
KOSELUGO ORAL CAPSULE 10	MAINT)	. 310
MG, 25 MG309	MAYZENT STARTER(FOR 2MG	
KRAZATI9	MAINT)	. 310
KYNMOBI SUBLINGUAL FILM 10	megestrol oral suspension 400 mg/10 ml	
MG, 10-15-20-25-30 MG, 15 MG, 20	(40 mg/ml)	.158
MG, 25 MG, 30 MG	megestrol oral tablet	
lanreotide	MEKINIST ORAL RECON SOLN	
<i>lapatinib</i> 183	MEKINIST ORAL TABLET 0.5 MG,	
lenalidomide	MG	
LENVIMA	MEKTOVI	
leuprolide (3 month)191	methocarbamol oral tablet 500 mg, 750	
leuprolide subcutaneous kit189	mg	152
lidocaine hcl mucous membrane solution 4	miglustat	
% (40 mg/ml)197	mimvey	
lidocaine topical adhesive patch,medicated	modafinil oral tablet 100 mg, 200 mg	
5 %197	morphine concentrate oral solution	
lidocaine topical ointment197	MVASI	
lidocaine-prilocaine topical cream 198	NATPARA	
LONSURF ORAL TABLET 15-6.14	NERLYNX	
MG, 20-8.19 MG	NEULASTA	
LORBRENA ORAL TABLET 100	NEULASTA ONPRO	
MG, 25 MG202	NINLARO	
LUMAKRAS ORAL TABLET 120	nitisinone	
MG, 320 MG324	NIVESTYM	
LUNSUMIO	NORDITROPIN FLEXPRO	
LUPRON DEPOT	norethindrone ac-eth estradiol oral tablet	
LUPRON DEPOT (3 MONTH) 193	0.5-2.5 mg-mcg, 1-5 mg-mcg	
LUPRON DEPOT (4 MONTH) 193	NOXAFIL ORAL SUSP, DELAYED	
LUPRON DEPOT (6 MONTH) 193	RELEASE FOR RECON	. 271
LYBALVI236	NUBEQA	
<i>lyllana</i> 148	NUCALA SUBCUTANEOUS AUTO	
LYNPARZA237	INJECTOR	
LYTGOBI	NUCALA SUBCUTANEOUS RECO	
MARGENZA206	SOLN	
MAVENCLAD (10 TABLET PACK) 64	NUCALA SUBCUTANEOUS	
MAVENCLAD (4 TABLET PACK) 64	SYRINGE 100 MG/ML, 40 MG/0.4	
MAVENCLAD (5 TABLET PACK)64	ML	.207
MAVENCLAD (6 TABLET PACK) 64	NUPLAZID	
MAVENCLAD (7 TABLET PACK) 64	NURTEC ODT	
MAVENCLAD (8 TABLET PACK)64	NYVEPRIA	
MAVENCLAD (9 TABLET PACK)64	OCALIVA	
MAYZENT ORAL TABLET 0.25 MG,	OCREVUS	
1 MG, 2 MG310	ODOMZO	
	OFEV	

OGIVRI356	posaconazole oral270	0
OJJAARA216	PREMARIN ORAL148	8
OLUMIANT32	PREMPHASE148	8
ONTRUZANT357	PREMPRO148	8
ONUREG30	PREVYMIS INTRAVENOUS	
OPDIVO231	SOLUTION 240 MG/12 ML, 480	
OPDUALAG232	MG/24 ML	8
OPSUMIT205	PREVYMIS ORAL188	8
ORENCIA3	PROMACTA ORAL POWDER IN	
ORENCIA (WITH MALTOSE)1	PACKET 12.5 MG, 25 MG104	4
ORENCIA CLICKJECT3	PROMACTA ORAL TABLET 12.5	
ORFADIN ORAL CAPSULE 20 MG. 230	MG, 25 MG, 50 MG, 75 MG 104	4
ORFADIN ORAL SUSPENSION 230	promethazine injection solution	2
ORGOVYX279	promethazine oral142	
ORILISSA ORAL TABLET 150 MG,	promethazine rectal142	
200 MG	promethegan rectal suppository 12.5 mg,	
ORKAMBI ORAL GRANULES IN	25 mg	2
PACKET203	pyrimethamine274	4
ORKAMBI ORAL TABLET203	QINLOCK	
ORSERDU ORAL TABLET 345 MG,	quinine sulfate27:	5
86 MG	QULIPTA20	
OTEZLA	RAVICTI138	
OTEZLA STARTER20	RELEUKO12:	5
OXLUMO204	RETACRIT INJECTION SOLUTION	
PALYNZIQ257	10,000 UNIT/ML, 2,000 UNIT/ML,	
paroxetine hcl oral suspension159	20,000 UNIT/2 ML, 20,000 UNIT/ML,	
paroxetine hcl oral tablet 159	3,000 UNIT/ML, 4,000 UNIT/ML,	
pazopanib249	40,000 UNIT/ML112	2
PEMAZYRE260	RETEVMO ORAL CAPSULE 40 MG,	
penicillamine oral tablet261	80 MG 308	8
phenobarbital160	REZLIDHIA238	8
PIQRAY ORAL TABLET 200	REZUROCK30	6
MG/DAY (200 MG X 1), 250 MG/DAY	RIABNI290	6
(200 MG X1-50 MG X1), 300 MG/DAY	RINVOQ360	6
(150 MG X 2)	RITUXAN HYCELA294	4
pirfenidone oral capsule265	ROZLYTREK ORAL CAPSULE 100	
pirfenidone oral tablet 267 mg, 534 mg,	MG, 200 MG109	9
<i>801 mg</i> 265	RUBRACA299	9
PLEGRIDY SUBCUTANEOUS PEN	RUXIENCE29	7
INJECTOR 125 MCG/0.5 ML, 63	RYBREVANT10	6
MCG/0.5 ML- 94 MCG/0.5 ML173	RYDAPT210	0
PLEGRIDY SUBCUTANEOUS	sajazir164	4
SYRINGE 125 MCG/0.5 ML, 63	sapropterin oral tablet, soluble30	1
MCG/0.5 ML- 94 MCG/0.5 ML173	SCEMBLIX22	
POMALYST	scopolamine base14	4

SEROSTIM SUBCUTANEOUS	TASIGNA ORAL CAPSULE 150 MG,
RECON SOLN 4 MG, 5 MG, 6 MG 320	200 MG, 50 MG225
SIGNIFOR248	tasimelteon
SIKLOS ORAL TABLET 100 MG 161	TAZVERIK331
sildenafil (pulm.hypertension) oral tablet 250	TECVAYLI334
SIRTURO34	TEPMETKO337
SKYRIZI291	teriflunomide338
<i>sodium oxybate</i> 312	testosterone cypionate340
SOMATULINE DEPOT	testosterone enanthate340
SUBCUTANEOUS SYRINGE 120	testosterone transdermal gel in metered-
MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3	dose pump 12.5 mg/ 1.25 gram (1 %),
ML182	20.25 mg/1.25 gram (1.62 %)
SOMAVERT258	testosterone transdermal gel in packet 1 %
sorafenib323	(25 mg/2.5gram), 1 % (50 mg/5 gram)340
SPRAVATO116	testosterone transdermal solution in
SPRYCEL ORAL TABLET 100 MG,	<i>metered pump w/app</i> 340
140 MG, 20 MG, 50 MG, 70 MG, 80	tetrabenazine341
MG78	THALOMID343
STELARA 369, 371	TIBSOVO177
STIVARGA278	TIVDAK344
STRENSIQ23	TRACLEER ORAL TABLET48
sunitinib malate326	TRACLEER ORAL TABLET FOR
SUNOSI	SUSPENSION48
SYMDEKO342	TRAZIMERA359
SYMLINPEN 120273	TRELSTAR INTRAMUSCULAR
SYMLINPEN 60273	SUSPENSION FOR
SYMPAZAN65	RECONSTITUTION363
SYNAREL218	TREMFYA140
SYNRIBO239	tretinoin352
TABRECTA58	trientine oral capsule 250 mg361
tadalafil (pulm. hypertension)250	TRIKAFTA ORAL TABLETS,
TAFINLAR ORAL CAPSULE72	SEQUENTIAL101
TAFINLAR ORAL TABLET FOR	TRUSELTIQ167
SUSPENSION	TRUXIMA295
TAGRISSO243	TUKYSA ORAL TABLET 150 MG, 50
TAKHZYRO SUBCUTANEOUS	MG364
SOLUTION181	TURALIO263
TAKHZYRO SUBCUTANEOUS	TYSABRI221
SYRINGE 150 MG/ML, 300 MG/2 ML	UBRELVY365
(150 MG/ML)181	UDENYCA254
TALTZ AUTOINJECTOR179	UDENYCA AUTOINJECTOR 254
TALTZ SYRINGE 179	UPTRAVI INTRAVENOUS305
TALVEY329	
TALZENNA	
TASCENSO ODT 129	

UPTRAVI ORAL TABLET 1,000	
MCG, 1,200 MCG, 1,400 MCG, 1,600	
MCG, 200 MCG, 400 MCG, 600 MCG	,
800 MCG	
UPTRAVI ORAL TABLETS, DOSE	
PACK	. 305
VANFLYTA	
VEGZELMA	42
VELCADE	
VENCLEXTA ORAL TABLET 10	
MG, 100 MG, 50 MG	.374
VENCLEXTA STARTING PACK	
VERZENIO	5
vigabatrin	
vigadrone	
VITRAKVI ORAL CAPSULE 100	
MG, 25 MG	.184
VITRAKVI ORAL SOLUTION	184
VIZIMPRO	
VONJO	
voriconazole oral suspension for	
reconstitution	.377
VOSEVI	
VOTRIENT	
VOWST	
VUMERITY	
WELIREG	
XALKORI	
XELJANZ	350
XELJANZ XR	350
XERMELO	
	84
XIFAXAN ORAL TABLET 200 MG,	
550 MG	.285
XOLAIR	
XOSPATA	
XPOVIO ORAL TABLET 100	
MG/WEEK (50 MG X 2), 40	
MG/WEEK (40 MG X 1), 40MG	
TWICE WEEK (40 MG X 2), 60	
MG/WEEK (60 MG X 1), 60MG	
TWICE WEEK (120 MG/WEEK), 80	
MG/WEEK (40 MG X 2), 80MG	
TWICE WEEK (160 MG/WEEK)	307
XTANDI ORAL CAPSULE	

XTANDI ORAL TABLET 40 MG, 80	
MG	. 110
XYOSTED	. 340
XYREM	
yargesa	213
YERVOY	
YONSA	
ZARXIO	
ZEJULA ORAL CAPSULE	
ZEJULA ORAL TABLET	228
ZELBORAF	373
ZIRABEV	44
ZOLADEX	
ZTALMY	
ZTLIDO	
ZYDELIG	
ZYKADIA	
ZYNLONTA	
ZYNYZ	